


## Act on Axial SpA Primary Care Presentation









### What is Axial Spondyloarthritis (Axial SpA)?

Umbrella term for inflammatory arthritis affecting spine and Sacroiliac joints.

- Ankylosing Spondylitis (AS) radiographic Axial SpA
  - Changes to the sacroiliac joints seen in x-ray
- Non radiographic Axial SpA (nr-AxSpA)
  - X-ray changes not present
  - Inflammation is visible on MRI
  - Person has a range of other symptoms











- Inflammatory pain
- Functional impairment
- Onset typically starts late teens early 20's (average age 26yrs)
- Can have lifelong impact and long term implications if left untreated
- Diagnosis is difficult and often delayed





## Axial SpA what are the key symptoms?



Inflammatory arthritis effecting spine and Sacroiliac joints.

- 1. Back pain started before the age of 40
- 2. Back pain developed slowly
- 3. Chronic back pain lasting greater than 3 months
- 4. Back stiffness upon waking
- 5. Back pain improves with exercise / movement
- 6. Back pain worse with rest
- 7. Alternating buttock pain
- 8. Regular waking at night

Symptoms starting slowly
Pain in the lower back
Improves with movement
Night time waking
Early onset (under 40)

Complete the NASS symptom checker\*: <u>Symptom checker</u>



### Gender in axial SpA

Axial SpA affects both women and men from a young age. It is no longer seen as a male disease. Females have a 2 year longer time to diagnosis around the

world.



- Men with axial SpA show a higher rate of radiological progression compared with women.
- Non-radiographic axial SpA (nr-axSpA) is more prevalent in women. (67% vs 33%)
- Ankylosing Spondylitis or radiographic axial SpA is more prevalent in men. (67% vs 33%)
- Women with axial SpA have, in general, higher disease activity scores (higher BASDAI, lower ASQoL, lower BASMI)
- Women with axial SpA also have more peripheral manifestations compared to males (enthesitis, IBD, Psoriasis and Peripheral Arthritis)

<sup>-</sup> Xabier Michelena, Clementina López-Medina, Helena Marzo-Ortega, Non-radiographic versus radiographic axSpA: what's in a name?, Rheumatology, Volume 59, Issue Supplement\_4, October 2020, Pages iv18-iv24, https://doi.org/10.1093/rheumatology/keaa422

<sup>-</sup> Boonen A et al. Semin Arthritis Rheum. 2015;44(5):556-562

<sup>-</sup> Rudwaleit M and Sieper J. Nat Rev Rheumatol. 2012;8(5):262-266

## Extra-musculoskeletal manifestations (EMM) of axial SpA?

Extra-musculoskeletal manifestations (EMMs) are common, important features of axial

spondyloarthritis (axial SpA).

The most prevalent being -

- acute anterior uveitis (AAU),
- inflammatory bowel disease (IBD) and
- psoriasis.

#### Other EMMs are:

- Enthesitis
- Dactylitis





## UK Economy: Paying a high price





Delay to diagnosis of axial SpA costs the UK economy

£18.7 billion

per year

**#WaitingCosts** 






Average time to diagnosis from symptom onset is 8.5 years



Waiting for an axial SpA diagnosis costs each person an average of £187k



#### One year costs less...

A Gold Standard time to diagnosis would save the UK economy

per person

Funded bu:












## Catalysing change with our health care ambassadors

#### Champions in Primary Care

Location of appointments for Champions in Primary Care programme







#### A2E Cohort 1

Current Cohort 1 participant rheumatology department / hospital



#### A2E Cohort 2

Current Cohort 2 participant rheumatology department / hospital



#### A2E Cohort 3

Cohort 3 participant rheumatology department / hospital

2 sites in cohort 2 & 1 cohort 3





## Our new Champions in Primary Care

- Shining a light on the issues facing achieving timely diagnosis in axial SpA
- Championing change and drives improvement in axial SpA care
- Sharing the story of change to showcase best practice improvements in axial SpA
- Spreading learning across healthcare systems to help other adopt change

#### **Champions in Primary Care programme**

A cadre of clinical champions in primary care and community services whose work ensures that axial spondyloarthritis (or axial SpA for short) is higher within the clinical reasoning of primary care professionals, so that patients who present with suspected axial SpA are identified at the first presentation and urgently referred to rheumatology.

Leadership Development Quality Improvement Training Local Improvement Projects

Action learning sets, Shared learning community, Celebrating Success



## Primary care – Barriers for diagnosis of Axial


SpA

- Lack of awareness about the disease amongst primary care workforce
- Non-specific findings on physical examination
- Diagnostic delay M-5.5 and F-8.8 NEI
- Lack of set diagnostic criteria
- Lack of biomarkers for diagnosis
- Lack of streamlined rheumatology pathways

Marina N et al 2020, Recognizing Axial Spondylarthritis: A Guide for Primary Care, A review, Mayo clinic

Crossfield S et al Changes in ankylosing spondylitis incidence, prevalence and time to diagnosis over two decades, BMJ, RMD open, Vol7 ,3

Russell M et al 2021, Diagnostic delay is common for patients with axial spondylarthritis: results from the National Early Inflammatory Arthritis Audit British Journal of Rheumatology , 61, 2



Marina N et al 2020, Recognizing Axial Spondylarthritis: A Guide for Primary Care, A review, Mayo clinic









- Create awareness of Axial SpA in primary care via education/learning modules
- Improve identification and screening in primary care
- Use of screening tools (SPADE/PRIMIS), guidelines (NICE/ASAS) and templates (ARDENS) to maximize the efficacy of patient consultations
- Improve referral pathways to Rheumatology
- Improve patient awareness / education and patient management in primary care
- Utilisation of best MSK practice guidelines for identification and referrals



#### How to find out more

• Go to <u>www.actonaxialspa.com</u> and read our first Act on Axial SpA campaign impact report.




www.nass.co.uk



## Resources for primary care HCPs



- NICE Guideline NG65
- GIRFT / BEST MSK pathway
- Ardens MSK Template
- SPADE Tool (1)
- Accurx Floreys
- Primis Pop up Tool
- ASAS / EULAR guidance











EUROPEAN ALLIANCE OF ASSOCIATIONS FOR RHEUMATOLOGY





## Back pain? Don't wait until it's too late.

Check the symptoms. It could be Axial SpA. See your GP.

**Get axial SpA** diagnosed.

Find out more at actonaxialspa.com







**Axial SpA** works silently. We don't.

Symptoms starting slowly Pain in the lower back Improves with movement Night time waking Early onset (under 40)



